



Nagoya City University Academic Repository

学位の種類	博士 (医学)
報告番号	甲第1690号
学位記番号	第1207号
氏名	森 賢人
授与年月日	平成 31年 3月 25日
学位論文の題名	Compensatory Increase in Heart Rate Is Responsible for Exercise Tolerance among Male Patients with Permanent Atrial Fibrillation (永続性心房細動男性症例の運動耐容能維持における運動時心拍数増加の意義) Tohoku J Exp Med 2018; 246: 265-274.
論文審査担当者	主査： 三島 晃 副査： 早野 順一郎, 大手 信之

Compensatory Increase in Heart Rate Is Responsible for Exercise Tolerance among Male Patients with Permanent Atrial Fibrillation

It is widely accepted that exercise tolerance in patients with atrial fibrillation (AF) is reduced because development of AF is accompanied by loss of atrial kick, resulting in decreased CO (Ostermaier et al. 1997). However, many patients with permanent AF, which lasts for at least a year without interruption, and preserved left ventricular ejection fraction (LVEF \geq 50%) are asymptomatic and have good exercise tolerance (Flaker et al. 2005; Myrstad et al. 2016). In such cases, the possible mechanism that compensates for the decrease in cardiac output accompanying the loss of atrial kick is a sufficient increase in heart rate (HR) during exercise. We investigated the relationship between exercise tolerance and peak HR during exercise using cardiopulmonary exercise testing in 242 male patients with preserved LVEF, 214 with sinus rhythm (SR) and 28 with permanent AF. Peak HR was significantly higher in the AF group than the SR group (148.9 ± 41.9 vs. 132.0 ± 22.0 beats/min, $p = 0.001$). However, oxygen uptake at peak exercise did not differ between the AF and SR groups (19.4 ± 5.7 vs. 21.6 ± 6.0 mL/kg/min, $p = 0.17$). In multiple regression analysis, peak HR (β , 0.091; $p < 0.001$) and the interaction term constructed by peak HR and presence of permanent AF (β , 0.05; $p = 0.04$) were selected as determinants for peak VO_2 ; however, presence of permanent AF was not selected (β , -0.38; $p = 0.31$). Therefore, the impact of peak HR on exercise tolerance differed between the AF and SR groups, suggesting that a sufficient increase in HR during exercise is an important factor to preserve exercise tolerance among patients with AF.