

Nagoya City University Academic Repository

学位の種類	博士(医学)
報告番号	甲第1427号
学位記番号	第1032号
氏 名	園田 浩生
授与年月日	平成 26年 3月 25日
学位論文の題名	Plasma N-Terminal Pro-Brain Natriuretic Peptide Levels Identifying Left Ventricular Diastolic Dysfunction in Patients With Preserved Ejection Fraction (左室拡張機能障害診断における NT-proBNP 測定の有用性) Circulation J 2012;76:2599-2605
論文審査担当者	主查: 三島 晃 副查: 早野 順一郎,大手 信之

Plasma N-terminal pro-brain natriuretic peptide levels identifying left ventricular diastolic dysfunction in patients with preserved ejection fraction.

Abstract

BACKGROUND:

Diagnosis of left ventricular (LV) diastolic dysfunction by blood testing is expedient in the clinical setting.

METHODS AND RESULTS:

In 98 patients with LV ejection fraction ≥50% who underwent cardiac catheterization for evaluation of coronary artery disease, LV pressure (LVP) was measured using a catheter-tipped micromanometer. A time constant, τ , of LV relaxation was computed from LVP decay; the inertia force of late systolic aortic flow, a surrogate index of LV elastic recoil, was also computed from the LVP-dP/dt relation (phase loop). Patients were classified into 2 groups: those with impaired LV relaxation (τ ≥48 ms) and those with preserved LV relaxation (τ <48 ms). Patients were also classified into another 2 groups: those with inertia force (≥0.5 mmHg) and those without (<0.5 mmHg). Plasma N-terminal pro-brain natriuretic peptide (NT-proBNP) ≥56.5 pg/ml had a sensitivity of 100%, specificity of 52.5%, and negative predictive value of 100% for identifying impaired LV relaxation. NT-proBNP ≥244.5 pg/ml had a sensitivity of 62.5% and specificity of 93.9% for detecting lack of inertia force.

CONCLUSIONS:

NT-proBNP level <56.5 pg/ml could be used as a value to sensitively identify patients with preserved LV systolic and diastolic function among those with coronary artery disease. NT-proBNP level ≥244.5 pg/ml is able to specifically detect a lack of inertia force and has potential for specifically diagnosing LV isolated diastolic dysfunction.