

Nagoya City University Academic Repository

学位の種類	博士(医学)
報告番号	甲第1670号
学位記番号	第1187号
氏 名	坂本 祐真
授与年月日	平成 31 年 3 月 25 日
学位論文の題名	CCR4 mutations associated with superior outcome of adult T-cell leukemia/lymphoma under mogamulizumab treatment (CCR4 遺伝子変異はモガムリズマブ (抗 CCR4 抗体) 治療における成人 T 細胞性白血病/リンパ腫の良好な予後と関連する) Blood 2018;132:758-761
論文審查担当者	主查: 山崎 小百合 副查: 高橋 智, 稲垣 宏

Title:

CCR4 mutations associated with superior outcome of adult T-cell leukemia/lymphoma under mogamulizumab treatment

ABSTRACT

Adult T-cell leukemia/lymphoma (ATL) has a dismal prognosis. CCR4 is expressed by tumor cells from most ATL patients, so therapeutic antibodies such as mogamulizumab, may be effective treatments. Here, we investigate whether gain-of-function mutations in the carboxyl terminus of CCR4, which were observed in 38 (33%) of 116 ATL patients, influence overall survival (OS) and response to treatment. We found no significant differences in OS when the whole patient cohort was stratified according to *CCR4* mutations. However, in those receiving mogamulizumab, 5-year survival from initiation of treatment in patients with (n = 11) or without (n = 31) *CCR4* mutations was 72.7% and 26.2%, respectively (P = 0.027). In contrast, *CCR4* mutations did not influence the outcome of allogeneic hematopoietic stem cell transplantation. Finally, in patients with aggressive-variant ATL, 5-year survival on mogamulizumab for patients with (n = 10) or without (n = 28) *CCR4* mutations was 80.0% and 24.7%, respectively (P = 0.006). These findings suggest that ATL patients with gain-offunction *CCR4* mutations are especially good responders to mogamulizumab-containing treatments. Thus, we conclude that *CCR4* gain-of-function mutations determine sensitivity to mogamulizumab therapy in ATL.